

# Package ‘ntranova’

April 10, 2024

**Title** Two Way Neutrosophic ANOVA

**Version** 0.0.1

**Description** Dealing with neutrosophic data of the form  $N=D+I$ (where  $N$  is a Neutrosophic number , $D$  is the determinant part of the number and  $I$  is the indeterminacy part) using the neutrosophic two way anova test keeps the type I error low. This algorithm calculates the fisher statistics when we have a neutrosophic data, also tests two hypothesizes, first is to test differences between treatments, and second is to test differences between sectors. For more information see Miari, Mahmoud; Anan, Mohamad Taher; Zeina, Mohamed Bisher(2022) <<https://www.americaspg.com/articleinfo/21/show/1058>>.

**License** GPL-3

**Encoding** UTF-8

**RoxygenNote** 7.3.1

**NeedsCompilation** no

**Author** Mohamad Taher Anan [aut, cre] (<<https://orcid.org/0009-0005-9468-6262>>),  
Mohamad Bisher Zeina [aut],  
Shaza Zubeadah [aut],  
Mahmoud Miari [aut]

**Maintainer** Mohamad Taher Anan <[mtanan200988@gmail.com](mailto:mtanan200988@gmail.com)>

**Repository** CRAN

**Date/Publication** 2024-04-10 17:00:02 UTC

## R topics documented:

ntaov	2
Index	3

---

ntaov

*Neutrosophic Two Way ANOVA*

---

**Description**

Neutrosophic Two Way ANOVA

**Usage**

```
ntaov(dt)
```

**Arguments**

dt is a data frame

**Value**

Neutrosophic ANOVA Table

**Examples**

```
y=c(4,5,3,9,11,8,15,12,14)
y1=c(6,7,5,11,14,10,17,13,16)
tr=c(1,1,1,2,2,2,3,3,3)
cek=c(1,2,3,1,2,3,1,2,3)
dt=data.frame(y,y1,tr,cek)
ntaov(dt)
```

# Index

ntaov, [2](#)